

LYSO-Easy

Lactic Acid Bacteria Inhibitor – Ready-To-Use Lysozyme Solution

LYSO-easy is a liquid solution of Lysozyme, ready to use. It has exactly the same application characteristics as powdered Lysozyme. 4 mL of Lyso-easy solution equals 1 gram of Lysozyme powder.

Lysozyme is a naturally occurring enzyme isolated from egg whites. It is used in wine to inhibit lactic acid bacteria. Lysozyme degrades the cell wall of gram-positive bacteria such as *Oenococcus*, *Pediococcus*, and *Lactobacillus*. Due to their protective external membranes, Lysozyme is not effective against gram-negative bacteria like *Acetobacter*. It also has no activity against yeast.

Lysozyme's effectiveness depends not only on the type of bacteria, but also the number of cells present. Unlike SO₂, Lysozyme is more effective at higher pH's when lactic acid bacteria growth is favored.

Lysozyme does not replace SO₂ because it has no anti-oxidative effect. It can, however, be used to help greatly reduce the amount of SO₂ needed to achieve microbial stability over the life of both red and white wines.

Lysozyme can be used in the following applications:

Lysozyme Applications	Reds	Whites	Recommended Dosage				Timing of Addition	
			Powdered Lysozyme		22% Lysozyme Solution			
Protection During Stuck and Sluggish Fermentations To encourage yeast growth in the absence of SO ₂ while reducing the risk of VA production by lactic acid bacteria.	•	•	250-400 ppm	25-40 g/hL	0.94-1.50 g/gal	114 - 182mL/hL	4.3 - 6.8 mL/gal	Add at first signs of a stuck fermentation
Prevent Growth of LAB in Must and Juice To inhibit spoilage characters due to uncontrolled microbial growth. This is especially important in high pH conditions or with grapes containing rot.	•	•	200 ppm	20 g/hL	0.75 g/gal	91.0 mL/hL	3.4 mL/gal	Add prior to fermentation
Delay MLF/Post-MLF Stabilization To protect wine without the negative effects of SO ₂ , to allow for maceration or aging, to allow for implantation of selected ML bacteria, or to increase efficiency of Phase I micro-oxygenation.	<i>Delay</i>	•	100-200 ppm	10-20 g/hL	0.38-0.75 g/gal	46 - 91 mL/hL	1.7 - 3.4 mL/gal	Add at juice stage or immediately after alcoholic fermentation
	<i>Stabilize</i>	•	250-500 ppm	25-50 g/hL	0.94-1.90 g/gal	114 - 228 mL/hL	4.3 - 8.6 mL/gal	Add immediately after MLF Completion
Inhibit MLF when Blending Partial and Complete ML Wines	•	•	300-500 ppm	30-50 g/hL	1.10-1.90 g/gal	136 - 227 mL/hL	5.0 - 8.6 mL/gal	Add during blending

Directions for Lab Scale Additions of LYSO-Easy

Prepare a 10% solution of LYSO-Easy by dissolving 10.0 ml of LYSO-Easy in 80 ml of cool or lukewarm water. When dissolved, bring volume up to 100 mL with DI water. Be sure to mix gently! Lysozyme is fragile and easily denatured by vigorous mixing or hot water.

For a 375ml Bottle:

<u>PPM Lysozyme</u>	<u>Milliliters of 10% LYSO-Easy Solution to Add</u>
100	1.73
150	2.56
200	3.41
250	4.28
300	5.10
400	6.83
500	8.55

 LALLEMAND

The information herein is true and accurate to the best of our knowledge; however, this data sheet is not to be considered as a guarantee expressed or implied or as a condition of sale of this product.

Document Edited 12/17/09